



COMBINING ABILITY STUDIES FOR YIELD AND YIELD COMPONENTS
IN SELECTED SOYBEAN LINES

Tadesse Ghiday^{1*}, Sentayehu A² and Asnake F³

¹Pawe Agricultural Research Center, Ethiopia

²College of Agriculture and Veterinary Medicine Jimma University

³Ethiopian Institute of Agricultural Research

ARTICLE INFO	ABSTRACT
Received 10 th April, 2016 Received in revised form 29 th May, 2016 Accepted 23 rd June, 2016 Published online 20 th July, 2016	In this study, 18 F ₁ hybrids obtained by crossing six lines (Ethiopia) with three testers (Brazil) in line × tester mating system during 2013/14 and were planted in randomized complete block design with two replications during 2014/15. Among the lines, BELESA 95 proved to be a good general combiner for number of pods per plant, grain yield and pod weight. Majority of the best specific combination for different characters resulted from the crosses among the parents with high x low and low x low GCA effects. Non-additive effects were predominant for number of pods per plant and grain yield, but appreciable additive effects were noted for hundred seed weight, pod weight and primary branches. The breeding method which can exploit non-additive as well as additive types of gene action is suggested for soybean improvement.
Keywords: Soybean, GCA, SCA	
<i>Copyright © 2016 Tadesse Ghiday, Sentayehu A and Asnake F., This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.</i>	

INTRODUCTION

In self pollinated crops like soybean (*Glycine max* (L.) Merrill), recombination breeding has been extensively used to develop the variability reservoir for exploitation in breeding program [1]. In a systematic breeding program, it is essential to identify the elite parents for hybridization, and superior crosses to expand the variability reservoirs for selection of superior genotypes [2]. Combining ability studies help in such endeavor [3]. In the present investigation, line x tester design with well adapted and widely grown varieties of soybean (tester) was used to obtain information on combining ability of elite lines for five characters of economic importance in soybean.

MATERIAL AND METHODS

Nine cultivar lines were selected on the basis of their geographical adaptation and morphological diversity. Out of them, three were testers (males), each crossed with six genotypes used as lines (females). All the testers (PSB2005-03, PSB2005-04 and PSB2005-06) were obtained from Brazil as introduced varieties (which are currently under production in Brazil). All the six lines (GIZO, AFGAT, GISHAMA, BELESA-95, ETHIOUGOSLVIA, and WOGAYEN) are high yielding Ethiopian varieties in the Southern West and Northern West parts of the country released for commercial production. Eighteen F₁s along with nine parents were planted in randomized complete block design with two replications

during 2014/2015 at experimental station of Pawe Agricultural Research Center. The parents were randomized among themselves. Each replication have single row of 5 meters length at 60cm x 10cm spacing. The mean data was recorded on ten random plants for five quantitative characters viz. number of pods per plant, grain yield (g), hundred seed weight (g), pod weight (g) and number of primary branches that was used for statistical analysis. The combining ability was done adapting Kempthorne [4] procedure. The total variance among F₁ hybrids was further partitioned into variance due to lines, testers and interaction component, which was used to estimated the additive and non-additive components of variance. Also, the contribution of lines, testers and their interaction towards total variability for each character was computed for assessing their relative importance according to Singh and Chaunhary [9].

RESULT AND DISCUSSION

Analysis of Variance

Analysis of variance indicated the presence of significant difference among the treatments for all the five characters studied (Table 1). The parents differed significantly for all the characters except grain yield; however, mean squares due to lines were significant only for hundred seed weight and pod weight. The testers differed significantly for number of pods per plant, grain yield and hundred seed weight. The hybrids showed significant differences only for pod weight. Further partitioning of variance among the hybrids showed that the mean square due

*✉ Corresponding author: Tadesse Ghiday
Pawe Agricultural Research Center, Ethiopia

to lines were significant for hundred seed weight and pod weight, due to testers for number of pods per plant, grain yield and hundred seed weight, and number of pods per plant in case of line x tester interactions (Table 1).

Table 1 Analysis of variance for five characters in Soybean at Pawe on station (2014/2015)

Source of variation	d.f	Number of pods per plant	Grain yield	Hundred seed weight	Pod weight	No. of primary branches
Replication	1	239**	63.3*	693.9**	0.05	41.1**
Treatments	26	71.7**	26.2*	127**	0.04**	3.3**
Parents	8	110.1**	24.4	307.1**	0.05**	8.0**
Parents vs crosses	1	190.4**	157.3**	0.3	0.1	0.1
Hybrids	17	50.8	21.9	55.6	0.03*	1.4
Lines	5	59.1	23.1	83.7*	0.07**	1.9
Testers	2	140.4*	63.6*	208.1**	0.02	1.1
Lines x testers	10	35.5**	16.1	22.5	0.02	1.1
error	26	29.3	15.1	43.6	0.02	1.7

significant at 5% and 1% levels, respectively

Table 2 Proportional contribution (%) to total variance for five characters in Soybean at Pawe on station (2014/2015)

Source of variation	Number of pods per plant	Grain yield (g)	Hundred seed weight (g)	Pod Weight (g)	No. of primary branches
Lines	35.8	32.5	46.3	63.7	43.6
Testers	21.3	22.4	28.8	5.6	6.1
Lines x testers	42.9	45.1	24.9	30.7	50.3

This was also clearly illustrated when the proportional contribution of each character studied. Lines and their interaction with testers contributed more than 70% of the total variance for all the characters. Except for number of pods per plant (21.3%), grain yield (22.4%) and hundred seed weight (28.8%), the contribution of testers was very little. The contribution of lines varied from 32.5% for grain yield to 63.7% for pod weight (Table 2).

Variance Components

The estimate of variance components (GCA and SCA) indicated that non-additive components were dominant for number of pods per plant and grain yield characters, though appreciable additive effects were noted for hundred seed weight, pod weight and primary branches (Table 3), as reported earlier by Mamta [5] and shanty [7]. These observations suggest that in soybean breeding, the methodology that can exploit both the additive as well as non-additive effects would be of immense value. Diallel selective mating of Shanti [6] and Sharma [5], which provides better opportunity for recombination, accumulation of desirable genes and selection would help in concentrating most of such genes in a pure line [11].

Table 3 Estimates of variance components for five characters in soybean at Pawe on station (2014/2015)

Source of variation	Number of pods per plant	Grain yield	Hundred seed weight	Pod weight	No. of primary branches
GCA	0.48	0.18	1.02	0.0001	0.08
SCA	3.07	0.47	-0.69	0.0001	-0.05
GCA/SCA	0.15	0.38	-1.47	1.000	-1.60

A judicious integration of the classical approach (Pedigree and

Bulk) with diallel selective mating may be of great help in achieving the quantum jump in soybean improvement [2] [8] and [10].

The estimates of GCA effects (Table 4) showed that among lines and testers, BELESA 95 was superior, as it showed positive and significant GCA effects for number of pods per plant and grain yield, beside pod weight. The lines ETHIOUGOSLAVIA and GIZO were good combiners for hundred seed weight (Table 4).

Table 4 General combining ability effect of the lines and testers for five characters in soybean at Pawe on station (2014/2015)

Source of variation	Number of pods per plant	Grain yield	Hundred seed weight	Pod weight	No. of primary branches
Lines					
AFGAT	2.47	0.89	0.57	0.08	0.41
WOGAYEN	1.11	0.48	0.55	-0.01	0.33
ETHIOUGOSLAVIA	-4.43	-1.67	4.27*	-0.03	0.93
GIZO	-0.49	-0.64	6.48*	-0.16*	-0.03
BELESA 95	5.89*	4.24*	2.00	0.11*	-0.37
GISHAMA	2.47	0.89	0.57	0.08	0.41
SE±	2.21	1.59	2.70	0.05	0.53
SE(gi-gj)	3.13	2.25	3.81	0.07	0.75
Testers					
PSB2005-03	1.48	1.30	3.07	-0.04	-0.24
PSB2005-06	1.74	0.86	0.59	0.03	-0.02
PSB2005-04	-3.22*	-2.16*	-3.66*	0.01	0.26
SE±	1.28	0.92	1.56	0.03	0.30
SE(gi-gj)	1.8	1.30	2.20	0.04	0.43

Significant at 5% level

Table 5 Specific combining ability of the best five crosses based on per se performances at Pawe on station (2014/2015)

Character	Cross	Mean	Sca Effect	Gca Status Of Parent	
				P1	P2
Number of pods per plant	BELESA 95 X PSB2005-03	44.2	6.57	High	Low
	ETHIOUGOSLAVIA X PSB2005-06	39.5	5.92	Low	Low
	BELESA 95 X PSB2005-06	37.2	-0.69	High	Low
	AFGAT X PSB2005-03	37.2	2.99	Low	Low
	GIZO X PSB2005-06	34.1	2.59	Low	Low
Grain yield (g)	BELESA 95 X PSB2005-03	30.0	4.82	High	Low
	ETHIOUGOSLAVIA X PSB2005-06	24.5	3.18	Low	Low
	BELESA 95 X PSB2005-06	24.3	2.42	High	Low
	AFGAT X PSB2005-03	22.4	-2.35	low	Low
	WOGAYEN X PSB2005-06	22.3	1.28	Low	Low
Hundred seeds weight (g)	WOGAYEN X PSB2005-03	44.5	4.48	Low	Low
	BELESA 95 X PSB2005-03	42.0	2.80	Low	High
	GIZO X PSB2005-03	41.1	-2.80	High	High
	GIZO X PSB2005-06	40.5	3.59	Low	Low
	GIZO X PSB2005-04	40.2	-1.12	Low	Low
Pod weight (g)	ETHIOUGOSLAVIA X PSB2005-06	1.1	0.07	Low	Low
	BELESA 95 X PSB2005-03	1.1	0.09	High	Low
	WOGAYEN X PSB2005-03	1.0	0.04	Low	Low
	AFGAT X PSB2005-06	1.0	0.03	Low	Low
	WOGAYEN X PSB2005-03	1.0	0.07	Low	Low
Primary branches	ETHIOUGOSLAVIA X PSB2005-03	7.3	1.32	Low	Low
	GISHAMA X PSB2005-03	7.3	0.75	Low	Low
	GIZO X PSB2005-04	7.2	1.18	Low	Low
	AFGAT X PSB2005-04	7.0	0.61	Low	Low
	WOGAYEN X PSB2005-04	6.7	-0.34	Low	Low

None of the testers was found to be a good general combiner. Association between per se performance and GCA effects was not evident in the present study. In fact, in many cases, the lines or testers with high mean had low GCA effects, indicating the ineffectiveness of choice of parents based on per se performance for hybridization (Table 5). Fourteen out of eighteen crosses occupied the first five ranks for five characters (Table 5). The four top ranking for grain yield proportionate to the order of number of pods per plants indicates the close association between number of pods per plant and grain yield. Of these fourteen crosses, eleven crosses were between low x low, two crosses between high x low and only one cross involved high x high gca parents. The largest number of low x low and high x low gca crosses in top ranks for different characters is of great interest, as such combinations could result in desirable transgressive segregants if the additive effects of one parent and the complementary epistatic effects (present in the cross) act in the same direction and maximize the expression of plant attributes under selection.

SUMMARY AND CONCLUSION

Two crosses Blessa95 x PSB2005-03 (high x low) and Ethiougoslavia x PSB2005-06 (low x low), exhibited high mean values for number of pods per plant and grain yield. These crosses may be further exploited for isolating the desirable segregates for economic traits of number of pods per plant and grain yield.

Acknowledgement

The authors greatly thank the Ethiopian Institute of Agricultural Research for financial support. Pawe Agricultural Research Center, staff members of the Soybean Research Team is greatly acknowledged for their active and unlimited cooperation in the execution of the experiment.

References

1. Bastawisy, M. H, Ibrahim, A. M and Mansours, S. H. 1997. Combining ability and heterosis studies for yield and its components in some top crosses of soybean [Glycine Max (L.)] Merril. *Annals of Agricultural Sciences, Moshtohor*. 35(1) : 93-106.
2. Darwish, I. H. I. 2007. Heterosis and inheritance of some quantitative character in soybean (Glycine Max L.). *Egyptian Journal of Plant Breeding*. 11(1) : 131-142.
3. El Sayad Z. S, Seliman, M. M.; Mokhtar, S. A, El Shaboury, H. M. G and ElHafez, G. A. A. 2005. Heterosis, combining ability and gene action in F₁ and F₂ diallel crosses among six soybean. *Genotypes*. 43(2) : 545-559.
4. Kamphrone, O. 1957. An introduction to genetic statistics. Johnwiley and Sons, Inc., New York. Kapila, R. K, Gupta, V. P and Rathore, P. K. 1994.
5. Mamta – Arya, Kamendra Singh, Pushendra and gupta, M. K. 2010. Heterosis study for yield, its components and quality characters in soybean [Glycine Max (L.) Merril]. *Soybean Research*. 8 : 75-79.
6. Shanti Patil, Vandana Khambalkar, Khedikar, Y. P and Menna Wankheda. 2003. Study of F₁ crosses of soybean. *Journal of Soils and Crops*. 13(1) : 112-115.
7. Shanti Patil; Vandana Kalamkar; Sable, N. H and Maheshwari, J. J. 2008. Study of heterosis in soybean. *Journal of Soil and Crops*. 18(1) : 208- 211.
8. Sharma, B, Singh, B. V, Kamendra Singh; Pushendra; Gupta, M. K. 2007. Selection criteria for improvement of grain yield in soybean [Glycine Max (L.) Merrill]. 5(1): 43-44.
9. Singh, R.K and Choudhary, B.D.1977. Line x Tester Analysis In : Biometrical methods in quantitative genetic analysis Kalyani Publishers, New Delhi. pp-178-85.
10. Sujata Bhat; Basavaraja, G. T and Salimath, P. M. 2011. Studies on genetic variability in segregating generation of soybean [Glycine Max (L.) Merril]. *Crop Research, Hisar*. 42(1/2/3) : 251-254.
11. Yang – Jia Yin and Gai Jun Yi. 2009a. Heterosis, combining ability and thin genetic basis of yield among key parental materials of soybean in Huang – Huai valleys. *Acta. Agronomica Sinica*. 35(4) : 620-630.
